

## **KINGSTON GENERAL HOSPITAL**

# **MEDICATION ADMINISTRATION: BELOW THE DRIP CHAMBER**

## **LEARNING GUIDE FOR REGISTERED NURSES AND REGISTERED PRACTICAL NURSES**

Prepared by: **Nursing Education**  
Date: **2001 November**  
Revised: **2002 March**  
Revised: **2010 November**

---

---

## TABLE OF CONTENTS

	<b>Page</b>
1.0 Introduction .....	3
1.1 Authorization Process .....	3
1.2 Expected Competencies for the Learner .....	3
2.0 Steps in the Administration of Medication Below the Drip Chamber .....	4
2.1 Medication Compatibility .....	4
2.2 Considerations Prior to Administration .....	5
2.3 Administration of Incompatible Medications .....	6
2.4 Administration of Compatible Medications .....	6
2.5 Common medications below the drip chamber.....	7
3.0 Respiratory Depression .....	7
3.1 Respiratory Depression .....	7
3.2 Naloxone .....	7
3.3 Documentation .....	8
4.0 Authorization Test: Adding Intravenous Medications Below the Drip Chamber .....	9
5.0 References .....	13
6.0 Evaluation of Learning Guide .....	14

Note: This learning guide contains information current at the time of distribution. Policies and procedures are frequently revised. Please refer to related policies and procedures contained in the Nursing Policy and Procedure Manuals for ongoing current information.

## 1.0 INTRODUCTION

There are three methods of administration for intravenous medications:

1. **Direct:**  
In direct administration, the medication is administered below the drip chamber, manually by syringe, either directly into the line or through an injection port over usually less than 10 minutes.
2. **Chamber:**  
The medication is administered above the drip chamber via minibag or Buretrol usually over 10 - 60 minutes.
3. **Infusion:**  
The medication is administered above the drip chamber over at least 60 minutes.

To avoid the potential complications of the administration of medications by the direct intravenous method, additional education and training is required. The administration of any intravenous medication below the drip chamber, manually by syringe over usually less than 10 minutes, is an **Added Nursing Skill for Registered Nurses (RN) and Registered Practical Nurses (RPN) and requires special authorization to perform.**

### 1.1 Authorization Process

To become authorized to administer medications below the drip chamber, the nurse will:

- review the learning guide related to administration of medications below the drip chamber;
- complete a written test with a passing level of 80%; and
- perform a minimum of one demonstration supervised by the Clinical Educator or delegate.

Reauthorization will be required when the need is identified by the Clinical Educator or delegate or by the nurse.

### 1.2 Expected Competencies for the Learner

Upon completion of the education program, the nurse will be able to:

- Administer an intravenous medication below the drip chamber.

## 2.0 STEPS IN THE ADMINISTRATION OF MEDICATION BELOW THE DRIP CHAMBER

Drugs that can be administered below the drip chamber are listed in the various drug references endorsed at KGH.

1. Become familiar with the properties of the drug that you are administering. Refer to the following endorsed references:
  - HDH/KGH Parenteral Drug Therapy Manual;
  - The KGH NICU Drug Manual will be used as the reference manual for neonates only in ICU, NICU and Pediatrics;
  - KGH/HDH Department of Pharmacy Services: Guidelines for the Admixture and Administration of Frequently Used IV Medications to Pediatric Patients will be used as a reference for infants and children;
  - The Hospital for Sick Children Handbook for Pediatrics is available as a reference for drug dosing for infants and children (located on the pediatric resuscitation cart in ER);
  - Computer program available for Pediatric medication dose calculation (ER, ICU, Peds);
  - The Hospital for Sick Children Formulary.
  
2. Select an Intravenous (IV) line. Consider:
  - what is infusing in the line (medications, blood products);
  - location and type of the intravenous line:

**Peripheral** lines may be prone to increased incidence of tissue damage. There may be more likelihood of decreased circulation in the vasoconstricted patient and therefore reduced distribution of the medication.

**Central** lines allow increased blood circulation. This allows for higher concentrations of the drug to be delivered. There will thus be a shorter reaction time for both therapeutic and adverse effects.
  
3. Check the IV catheter site for patency noting any signs of infiltration.
  
4. Determine compatibility of the medication with the IV solution. Refer to the HDH/KGH Parenteral Drug Therapy Manual and/or IV Drug Administration information.

### 2.1 Medication Compatibility

#### Types of Incompatibility

- Physical insolubility - An insoluble reaction results in a visible precipitate.
- Physical - There is no physical visible change, but the drug is affected and is less active.
- Chemical reaction - A reaction such as an acid-base reaction, can lead to precipitation, colour changes, gas

solution formation, or no observable change.

- pH incompatibility                      - The pH of the drug plus another drug or solution may lead to a reaction;
- Therapeutic interaction                - An interaction occurs within the body that may inactivate or potentiate the drug.

## **2.2 Considerations Prior to Administration**

Check: correct patient, route, dose, drug and time.

Before IV medication administration, consider:

- IV solution;
- pH of solution and drug;
- other drugs infusing in the IV line;
- the maximum concentration and rate of administration;
- drugs and solutions infused through a buretrol may remain in the filter, therefore, always flush the Buretrol;
- policy and procedure requirements;
- is the presence of a physician at the bedside or on the unit required?
- the patient condition and expected response to the medication
- If the patient is unstable or in an acute episode of illness the care of the patient and/or any drug administration should be an RN responsibility i.e. the patient in severe hypoglycemia should not receive Dextrose IV by an RPN

Due to the nature of some drugs, there may be the need for additional support and monitoring during and/or following their IV administration. These support and monitoring prerequisites are included in the Administration section of the relevant drug monographs of the HDH/KGH Parenteral Drug Therapy Manual. In order to highlight the drugs that have specific prerequisites, they also appear in a text box on the upper right hand corner of these drug monographs. The administration guidelines of all drugs should also be referred to in order to determine if there are other considerations in addition to the support and monitoring requirements. The support and monitoring requirements are defined as follows:

### **Blood Pressure Monitoring**

Drugs requiring the use of non-invasive blood pressure monitoring with vital signs being monitored more frequently than q1h,

**OR**

Those drugs requiring continuous arterial line blood pressure monitoring.

### **Cardiac Monitoring**

The patient must be connected to a cardiac monitor (excluding telemetry) and the Regulated Health Professional administering the drug must continuously monitor the patient and have acquired the skill of ECG interpretation.

### **Fetal Monitoring**

The patient must be connected to an electronic fetal monitor and the Regulated Health Professional must have acquired the skill of interpreting the monitor strip.

### **Respiratory Support**

There must be an oral airway, manual ventilator (ambu bag), airway suctioning and oxygen equipment readily available on the unit.

### **Ventilator Support**

The patient must have an endotracheal tube or tracheostomy tube in situ, and must be mechanically ventilated.

## **2.3 Administration of Incompatible Medications**

To administer incompatible medications:

1. Determine that the IV is patent.
2. Pinch/clamp the IV line just above the closest port to the patient, therefore preventing backflow during direct IV push administration.
3. Insert a 10cc syringe of compatible flush solution and flush the tubing with 10mL of flush, from the port to the patient.
4. Insert the syringe with medication and administer the medication according to dosage and rate guidelines.
5. Insert a 10cc syringe of compatible flush solution and flush the tubing with 10mL of solution from the port to the patient at a rate equal to the recommended rate of administration of the drug.
6. Resume the IV infusion as prescribed or flush saline lock as per policy I-5600. Document on the Medication Administration Record (MAR) or unit-specific flow sheet.

## **2.4 Administration of Compatible Medications**

1. Pinch the IV line just above the closest port to the patient.
2. Insert the syringe with the medication and administer according to dosage and rate guidelines. You may alternate pinching and pushing with releasing and infusing IV solution to meet the required rate guidelines.
3. Resume the IV infusion as prescribed or flush saline lock as per policy I-5600. Document on the Medication Administration Record (MAR) or unit-specific flow sheet.

## **2.5 Common Medications that can be Administered Below the Drip Chamber**

Heparin, decadron, nalaxone, dextrose, and gravol to name a few may be used on units throughout the hospital but always refer to the HDH/KGH Parenteral Manual to seek direction regarding the time frame which is recommended for drug administration. If the medication requires a physician present, specific monitoring requirements or if you are unsure in any way it should not be administered below the drip chamber.

## **3.0 RESPIRATORY DEPRESSION**

### **3.1 Respiratory Depression**

Respiratory depression caused by opioids is a decrease in the depth or rate of respirations per minute. Opioid related respiratory depression is more likely to occur if:

- the patient is >70 and receives a large dose;
- there is a history of impaired respiratory function;
- parenteral narcotics, sedatives or antiemetics were administered concurrently;
- drugs were administered during surgery and/or labour.
- patient has been lying flat;
- patient is obese.

Usually opioids produce sedation before producing respiratory depression. Therefore monitoring the level of sedation (LOS) is as important as checking the respiratory rate (Pastero & McCaffery 1994).

#### **Treatment**

- Notify the attending service if an adult or pediatric RR < 10 and/or decreased level of sedation (LOS).
- Administer oxygen.
- Administer naloxone IV as ordered.

### **3.2 Naloxone**

Naloxone is a pure opiate antagonist that rapidly reverses the central nervous system depression, respiratory depression and analgesic effect caused by opiate analgesics such as fentanyl. Naloxone should always be diluted and given in small increments.

### **Important Points to Remember**

- Be aware that as naloxone reverses the analgesic effect of an opioid, effective pain control becomes a concern.
- Continue symptomatic treatment of respiratory depression with oxygen and artificial ventilation (with oral airway and ambu bag) as necessary until naloxone is effective.
- When an excessive dose of naloxone has been used it can precipitate withdrawal syndrome in an opioid-dependent individual.
- Rapid Naloxone administration increases the risk of cardiovascular complications (CPS 2000).
- The RPN should not administer below the drip chamber.

### **3.3 Documentation**

Record the patient's assessment data, reaction to the drug, its effectiveness, and any adverse effects noted.



#### **4.0 AUTHORIZATION TEST: ADDING INTRAVENOUS MEDICATIONS BELOW THE DRIP CHAMBER**

On the answer sheet following the test questions, circle the letter representing the phrase that best completes the following statements.

1. Which of the following should be considered when selecting an IV line for direct medication administration?
  - a. solution currently infusing in the line
  - b. peripheral versus central line
  - c. maximum concentration and rate of administration
  - d. all of the above
  
2. What are the disadvantages of a peripheral IV line when compared to a central venous line for direct IV medication administration?
  1. increased incidence of tissue damage
  2. onset of adverse reactions is unchanged
  3. decreased blood circulation
  4. greater risk of air emboli
  - a. 1, 2 & 3
  - b. 2 & 4
  - c. 1 & 3
  - d. 1, 2, 3 & 4
  
3. Prior to administering a direct IV medication, the nurse must always ensure that the:
  1. catheter site shows no sign of infiltration
  2. IV line is patent and IV fluid is infusing well
  3. IV solution is compatible with the medication to be administered
  4. patient is not currently receiving a medication that is incompatible with direct IV medication
  5. dosage and rate of administration is correct
  - a. 1, 2, 3
  - b. 1, 2 & 5
  - c. 2, 3 & 4
  - d. all of the above

4. Which is the correct order of the following steps in the direct IV administration of a compatible medication:
1. Administer medication according to dosage and rate guidelines.
  2. Pinch the IV line above the closest port to the patient, swab port with alcohol swab and insert syringe with medication.
  3. Observe the patient for therapeutic or adverse reactions and complete the documentation.
  4. Remove syringe, recheck IV infusion rate.
  5. Check for infiltration and patency, ensuring an adequate infusion rate.
- a. 5, 2, 3, 1, 4  
b. 2, 5, 3, 1, 4  
c. 2, 1, 4, 3, 5  
d. 5, 2, 1, 4, 3
5. When administering a direct IV medication into an IV line with an incompatible solution, the nurse should
1. flush the line with 10cc of a compatible IV solution prior to administration of the drug
  2. administer the medication according to the dosage and rate guidelines
  3. alternate pinching of the IV line and pushing of the medication with releasing of the IV line and infusing the IV solution
  4. flush the IV line with 10cc of a compatible IV solution after administration of the drug at the same rate that the drug was administered
- a. 1 & 4  
b. 1, 2 & 4  
c. 1 & 2  
d. all of the above
6. Naloxone will reverse respiratory depression caused by the following medications:
1. morphine
  2. diazepam
  3. meperidine
  4. phenobarbital
- a. 1 & 2  
b. 1 & 3  
c. 2 & 4  
d. 3 & 4

7. If you are asked to give Diazepam 10mg parenterally how could you determine the time required to administer the drug and whether it is safe to give it below the drip chamber?
  - a. Request the doctor order the time frame with the dose
  - b. Call the pharmacist
  - c. Refer to the HDH/KGH parenteral drug therapy manual
  - d. Ask a nursing colleague

Indicate on the answer sheet whether the following statements are true (T) or false (F).

8. It is within the RPN scope of practice to give dextrose IV below the drip chamber to his/her patient that is experiencing severe hypoglycemia.
9. When administering medications that are incompatible, the typical flush solution is 10mL normal saline.
10. A chemical reaction of incompatible solutions will always lead to a visible change in the solution to be administered.

**Authorization for Administration of Drugs Below the Drip Chamber: Test Answer Sheet**

**Name** \_\_\_\_\_ **Date** \_\_\_\_\_ **Score** \_\_\_\_\_

- 1. a b c d
- 2. a b c d
- 3. a b c d
- 4. a b c d
- 5. a b c d
- 6. a b c d
- 7. a b c d
- 8. T F
- 9. T T
- 10. T F

## 5.0 REFERENCES

Compendium of pharmaceuticals and specialties (2000) 35<sup>th</sup> ed. Ottawa ON: Canadian Pharmaceutical Association.

Feenstra, G., & Connell, K. (2000). Learning module for direct IV administration of naloxone. Halifax, NS: Queen Elizabeth II Health Sciences Centre

Hotel Dieu Hospital/Kingston General Hospital Parenteral Drug Therapy Manual.  
Kingston ON:

Kingston General Hospital Nursing Policy & Procedure Manual. M-1670.

McCaffery, M., & Pasero, C. (1999). Pain clinical manual. 2<sup>nd</sup> ed. St. Louis MO: Mosby.

## 6.0 EVALUATION OF LEARNING GUIDE

Your feedback and comments are most appreciated. Thank you for your time in responding to this questionnaire. Your response will help us in planning/revising learning materials.

Please circle the appropriate response

Strongly disagree

Strongly agree

1. The content was clear and easy to understand.  
Comments:

1      2      3      4      5

2. The content was relevant.  
Comments:

1      2      3      4      5

3. I feel that my learning needs were met.  
Comments:

1      2      3      4      5

4. This guide will help me to meet the knowledge/skill requirements of administration of IV medications below the drip chamber.  
Comments:

1      2      3      4      5

Additional comments/suggestions:

Please return the completed evaluation to your Clinical Educator. Thank you.