

A New Beginning...Colostomy Care

Information for patients who are preparing
to receive a colostomy (or who have a colostomy).



Hôpital
Hotel Dieu
Hospital



Hôpital Général de
Kingston General
Hospital

Contents

Introduction	4
The digestive tract	5
What is a colostomy?	6
Why you need a colostomy	9
Before your operation	9
During your operation	11
After your operation	12
About your stoma	14
Note your observations:	14
About your pouching system	14
Pouch care	17
When to empty your drainable pouch	17
How to empty and clean your drainable pouch	18
How to remove your drainable pouch (2-piece system only)	18
Changing your pouching system	19
When to change your pouching system	19
How to change your pouching system	19
Note the features you prefer about pouching systems:	21
What about food?	22
Excessive gas	22
Odour	23
Diarrhea	24
Constipation	25
What to wear?	25
How to bathe and shower	26
What about work?	26
What about travel?	27

What about physical activity?27

 How to get out of bed after surgery28

 Temporary physical limitations28

 Returning/starting a new sport.....29

 Swimming and other water activities29

 Hernia prevention30

What about intimacy?.....30

Medications31

Adjusting to your colostomy31

Before you go home32

Post-surgical follow-up32

How to pay for supplies33

What to do if.....34

 Note your stoma/skin/pouching system concerns35

About your colostomy closure36

Information to remember37

Ostomy supplies list38

Helpful resources39

 Additional resources40

Glossary41

Notes43



Introduction


Your surgeon has told you that you need to have a colostomy. You may have a lot of questions, but cannot find the words to ask them. Many people feel this way before their colostomy surgery. You are not alone. Many people live long and full lives after their colostomy surgery. For them and for you, it is a new beginning.

This book is yours to keep. It tells you about your operation and the colostomy. This book also tells you how to care for your colostomy. It also tells you how to return to your usual way of life.

There is space for you to write notes or questions on page 43 to 44. Many health care providers are available to assist you and your family such as nurses, surgeons, social workers, physiotherapists, dietitians and pharmacists (and others) as you prepare and recover from surgery.

The digestive tract

The digestive tract begins at the mouth where the food is chewed and broken down. When swallowed, the food passes down the esophagus and enters the stomach. The food remains in the stomach for a short time as the acids and enzymes break down the food even further. The food matter then enters the bowel (or intestine). This is a tube-like structure that is divided into two main parts: small bowel (or small intestine) and large bowel (or large intestine). The small bowel absorbs the vitamins and minerals necessary for good nutrition. The small bowel has 3 parts: duodenum, jejunum and ileum. The large bowel (also called the colon) removes water and salts from waste that the body needs to keep. This waste is also known as stool. The waste changes from a liquid to a solid as it travels further down the large bowel. Muscles in the colon contract and relax to propel the stool along the length of the bowel into the rectum. Once the rectum is full, the muscles at the anus relax to allow for the stool to leave the body. This is referred to as a bowel movement.




What is a colostomy?

A colostomy is a surgical opening through the wall of the abdomen into the large bowel. The large bowel (colon) is brought out onto the surface of the skin on the abdomen to make an opening. This opening is called a stoma. With a colostomy, you pass stool through the stoma instead of storing it in the rectum and eliminating it through the anus.

There are four kinds of colostomies. Each one is named for the part of the large bowel where the colostomy is placed. You will have either an ascending, transverse, descending or sigmoid colostomy.

Anatomy of the Large Intestine



The frequency of the bowel movements and the consistency of the stool are different depending on the location of the colostomy.

The stool from an ascending or transverse colostomy is loose and watery. There may be three to four small bowel movements per day.


The stool from a descending or sigmoid colostomy is more formed. There may be one or two bowel movements per day.

Most commonly, the stoma is placed in the lower left side of your abdomen.




With a colostomy, you cannot control your bowel movements. You need to wear a pouching system over the stoma to collect the stool. The pouching system consists of a skin barrier and an odor-proof pouch.

The two most common types of colostomy construction are: end and loop.



End colostomy: The surgeon stitches the end of the large bowel to the abdomen. There is one opening into the colon. Stool and gas pass through this opening. The stoma is often on the lower left side of the abdomen.

End colostomy



Loop colostomy: The surgeon stitches a loop of large bowel to the abdomen. There are two openings in the stoma. One opening leads to the functioning part of the colon and will pass stool and gas. The second leads to the non-functioning part of the colon and allows mucus to pass through it.

Loop colostomy

Your colostomy may be temporary or permanent. This depends on the reason for your operation. A temporary colostomy is reversed after your bowel has healed. A permanent colostomy is not reversed. Your surgeon can tell you what type you will have and whether or not a portion of the colon and rectum were removed at the time of the operation.

Why you need a colostomy

A colostomy operation is done only if your health is at risk or if you can no longer cope with your condition.

There are many reasons for needing a colostomy. Your surgeon will explain why you need a colostomy and tells you what kind you will have.

Possible reasons for needing a colostomy may include:

- Colon, rectal or anal cancer
- Inflammatory bowel disease
- Diverticulitis
- Birth defects
- Trauma

Before your operation

Your surgeon tells you the date of your operation and arranges for you to come to the Pre-Surgical Screening Centre for tests before your operation. This may include x-rays, a cardiogram, and blood tests. It is always possible that your surgery may be cancelled due to an unexpected emergency.

Kingston Health Sciences Centre has Nurses Specialized in Wound Ostomy Continence, referred to as the NSWOC nurses. They help you learn about your colostomy. For planned surgeries, the NSWOC nurse from Kingston General Hospital site (KGH) meets with you to explain what to expect at the time of your operation. The NSWOC nurse will discuss the impact of having an ostomy on your day-to-day activities and will teach you how to take care of your ostomy. Potential complications and how to manage them are also discussed.

In addition, the NSWOC nurse examines your abdomen before surgery and selects the best location for your stoma. Your input is helpful since your lifestyle, clothing and specific needs are considered before choosing the site. Every effort is made to place the stoma in the best possible location. Sites vary depending upon the type of surgery and the anatomy of each person.

Your surgeon tells you how to prepare your bowels for your operation and may arrange for you to take a laxative. This is to make sure your bowels are empty before the operation. You may pass several watery stools after taking the laxative. Many colostomy surgeries are done on an urgent or emergency basis. In emergency cases, these preparations are not possible.

You may eat solid foods until 12 midnight the night before your surgery unless you had a bowel prep (laxative). If you had a bowel prep, follow the instructions that were given to you.

You may have up to 3 glasses (800mL) of a high carbohydrate (sugary) drink the night before surgery, and 1.5 glasses (400mL) carbohydrate rich drink 3 hours before surgery. It is important to have sugary drinks before your surgery because it will help you feel stronger after your surgery and recover faster. Examples of high carbohydrate drinks are: apple juice, cranberry cocktail, and iced tea. **Do not eat or drink anything within 3 hours of your surgery.**

On the morning of your operation, you will register in the Same Day Admission Centre at KGH which is located on the second floor of the Connell Wing. After you've registered, you will then meet your nurse who will ask you a few questions. Your nurse prepares you for your operation. You put on a hospital gown. An intravenous (IV) is placed in your arm. This gives you fluids and medications that you need during the operation. Once this process is complete, you will be transferred to the operating room. If your family will be staying in the hospital while you are in surgery, they'll be shown to the waiting room in the Same Day Admission Centre.



During your operation

Your surgeon performs your operation through an incision in your abdomen.

Your surgery can be done in 3 ways:

- **Laparoscopic:** If your surgery will be done laparoscopically, the surgeon will make 4 to 6 small cuts (incisions) in your abdomen. Your surgeon will use long-handled instruments and a camera to loosen (free-up) the diseased bowel and then remove the section of bowel through a small incision. The surgeon can view images projected from inside the body onto a video screen. The healthy ends of your bowel will then be sewn or stapled back together and a stoma will be created.
- **Robotic:** If your surgery will be done robotically, the surgeon will make 4 to 6 small cuts (incisions) in your abdomen. Your surgeon will sit at a console beside you to operate tiny instruments that can rotate with more precision than a laparoscopy and a camera. The instruments will loosen (free-up) the diseased bowel and then remove the section of bowel through a small incision. With the camera, the surgeon can view high definition, 3-D magnified images from inside your body on the console screen. The healthy ends of you bowel will then be sewn or stapled back together and a stoma will be created.
- **Open:** If the surgery is done with a traditional open approach, your surgeon will make one 10 to 20cm cut (incision) in your abdomen to perform the surgery. The surgeon can see directly inside your body. The surgeon will remove the diseased section of bowel and sew the healthy ends of your bowel back together and a stoma will be created.

You may require the following:

- **Abdominal or pelvic drains.** These small tubes drain away extra fluid from your abdomen and pelvis.
- **Support stockings or special inflatable stockings.** These help the circulation in your legs and prevent blood clots.
- **Urinary catheter.** This tube goes into your bladder. It drains urine from your bladder until you are moving well enough to get up to pass urine on your own.

You will require the following:

- A **pouching system** is placed over your colostomy.
- A **dressings** is placed over your incision(s).

This will happen while you are still asleep during the surgery.

After your operation

After your operation, you are taken to the recovery room until you are awake. Then you go to a surgical unit. Your nurse:

- Makes sure that you are comfortable
- Checks your tubes, drains and vital signs
- Looks at your incision and colostomy
- Helps you with your breathing and leg exercises

It is important to remember everybody's experience may be different, but this is what you might expect:

Day/Evening of surgery:

- You may start drinking clear fluids 2 hours after your surgery
- You sit up at the side of the bed and dangle your legs

The first day after surgery:

- Chew 1 piece of gum 3 times a day for up to 5 minutes
- You may start to introduce solid foods
- You sit up in a chair for all of your meals or for at least 1 hour
- You may walk in the hallway as tolerated
- Empty your pouching system with a nurse

In 2 to 3 days:

- Chew 1 piece of gum 3 times a day for up to 5 minutes
- Eat solid food as tolerated or as ordered by your doctor
- Sit up in a chair for all meals
- You may walk in the hallway 4 times per day, as tolerated
- Empty your pouching system on your own
- Participate in pouching system change

In 3-4 days:

- Chew 1 piece of gum 3 times a day for up to 5 minutes
- Eat solid food as tolerated or as ordered by your doctor
- Sit up in a chair for all meals
- You may walk in the hallway a minimum of 4 times per day, as tolerated
- Fully participate in pouching system change
- Discuss and understand any dietary changes
- Review information about potential complications, products and financial supports
- Be discharged from hospital

Your colostomy may not function for a few days after your surgery. At first, you may have abdominal cramps and bloating. Then, gas starts to pass through the stoma. This means that your small bowel is beginning to work. Later, your colostomy passes liquid stool. As you return to a regular diet, there will be less gas and the stool becomes more formed.

After your operation, the NSWOC nurse fits you with a new pouching system. As you gradually recover from your surgery, the NSWOC nurse teaches you how to take care of your colostomy and prepares you for going home. It is important that you become as independent and comfortable as possible with caring for your colostomy. The floor nurses will also reinforce the teaching and support you in your learning. With time, you will develop confidence and the care becomes routine to you. Support from your family and friends is very helpful. If you wish, someone close to you may be included in your teaching sessions. Never hesitate to ask questions.

Prior to your discharge, your surgeon wants to ensure you have met three milestones which are:

- Being able to tolerate your food
- Having a return of bowel function
- Having your pain under control

About your stoma

The visible part of the colostomy is the stoma. This is the inner lining of the small bowel and it soft, moist and red in colour. At first, you may not feel ready to look at your stoma.

- There are no nerve endings in the stoma, so you have no feeling in it. It does not hurt, and you cannot tell when stool or gas is passing through it.
- The stoma may be oval or round. At first, it is a little swollen. The swelling goes away in 6-8 weeks.
- A clear gel covers your stoma. This is mucus that coats the inside of the bowel.
- The stoma bleeds easily since there are many small blood vessels close to the surface. The pouching system protects it well in case it is bumped.
- The stitches around the stoma dissolve in 2 to 3 weeks.
- The skin around your stoma may be bruised from the operation. The bruising lasts 1 to 2 weeks. It may last longer if you take drugs that thin your blood.

Note your observations:

About your pouching system

The colostomy pouching system is a system made to contain the stool and gas from your colostomy. It is made up of a skin barrier to protect the skin around your colostomy and an odour-proof pouch. Pouching systems are also lightweight, waterproof, and low profile (not noticeable under clothing).

There are a wide variety of pouching systems available. Your choice depends on your body type, personal preference and your lifestyle. The NSWOC nurse shows you pouch samples and helps you with your selection.

There are generally two kinds of pouching systems: one and two-piece systems.



The **1-piece pouching system**: The skin barrier and the pouch are pre-attached.



The **2-piece pouching system**: The pouch can be removed from the skin barrier. The skin barrier has a plastic ring. The pouch snaps onto or sticks to the plastic ring depending on the manufacturer.

There are also 2 kinds of pouches: **drainable** and **closed end**. These are also available in clear or opaque colours. You will have a clear pouch while you are in hospital to allow for viewing of your stoma by your surgeon and your nurse. You can change to an opaque pouch after you are discharged if you prefer.



Drainable pouch: The bottom of the pouch has an opening so you can empty it. The opening is closed with a clamp or other type of closure system.




Closed end pouch: There is no opening at the lower end of the pouch. You change the pouch after each bowel movement. This kind of pouch is also known as a non-drainable or disposable pouch. Closed end pouches come in different sizes. Small closed end pouches are ideal for intimacy or when swimming. These pouches have a limited capacity and may not be practical if you have diarrhea during an illness.

Pouch filter

A pouch filter prevents gas from inflating your pouch. The NSWOC nurse shows you how to add a filter to your pouch. You can also obtain a pouch with a built-in filter (see photo above).

The NSWOC nurse will tell you if you need to use an ostomy belt to secure your pouching system.



The belt fastens onto the tabs of the pouching system and must be worn at the level of the stoma. The NSWOC nurse will show you how to adjust the belt.

Pouch care

After your surgery, you will have a 2-piece pouching system with a drainable pouch. This is because your bowel movements are watery and your pouch is emptied often.

Learning how to care for your colostomy will be new for you. If you wish, you can have a family member or friend with you. It is important you become as independent and comfortable as possible with the care and management of your colostomy.

When to empty your drainable pouch

- When the pouch is a third full
- Before going out
- Before going to bed
- Before physical activity or exercise
- Before sexual activity
- Whenever it is convenient for you

How to empty and clean your drainable pouch

1. Sit well back on the toilet so you can empty the pouch into the toilet between your legs.
If this is not comfortable, you can:
 - put a chair beside the toilet and bend over the toilet
 - stand and empty the pouch into a container which you will later empty into the toilet.
2. Put a few sheets of toilet paper on top of the water in the toilet. This helps to prevent splashing.
3. Hold the end of the pouch up and remove the clamp.
4. Pinch the end of the pouch closed with your fingers.
5. Direct the end of the pouch into the toilet (or container) and release your pinched fingers to empty the contents.
6. Fill a plastic squeeze bottle with tap water. You may also add 5 to 10ml (1 to 2 teaspoons) of mouthwash. This helps control odour.
7. Squirt water into the pouch and move it around to clean the inside of the pouch. Empty the contents into the toilet. Repeat this until the pouch is clean.
8. Hold the pouch end in one hand, and wipe the outside and the bottom 2" inside the pouch with toilet paper.
9. Replace the clamp or close the end. Tug it slightly to make sure it is closed.

How to remove your drainable pouch (2-piece system only)

You may choose to remove the pouch that must be emptied and snap a clean pouch onto the skin barrier. The soiled pouch can then be emptied and rinsed. Two pouches can be alternated in this way for 1 to 2 weeks.

1. Hold the skin barrier against your abdomen. At the same time, gently pull the tab on the top of the pouch until the pouch comes off.
2. Wipe your stoma and inside the flange with toilet paper.
3. Attach a clean pouch. Starting at the bottom of the pouching system, line up the plastic rings. At the same time, press the pouch against the ring. It should snap (or stick) into place and feel secure.
4. Hold the skin barrier against your abdomen and gently tug the pouch downward. This tells you if it is secure.

5. Replace the clamp or close the end.
6. Empty the soiled pouch into the toilet.
7. Hold the pouch under the tap and run cool water through it until it is clean.
8. Pat the pouch dry with paper towel.

Changing your pouching system

The wear time of your pouching system depends on the amount and character of your stool. It is also dependent on the climate in which you live, your daily activities and the type of skin barrier you use. There are two types of skin barriers: standard and long-wear. The NSWOC nurse helps you choose the type that best suits your needs.

When to change your pouching system

- At least every 5 to 7 days
- When it is leaking
- When your skin under the skin barrier burns or feels itchy

It is best to change your pouching system before a meal, or at least two hours after a meal, when your colostomy is less active. A good time may be before breakfast or bedtime. When your incision is completely healed, you may take a shower or bath when the pouching system is off.

How to change your pouching system

1. Pull off the pouching system gently with one hand. At the same time, hold your skin back with the other hand. Discard the used pouching system in the garbage. **Do not flush it down the toilet.**
2. Wash the stoma and skin with warm water and a soft wash cloth. If you use soap, choose a soap that does not contain oil. Oil prevents the pouching system from sticking to your skin. **Many commercial skin wipes such as baby wipes contain lanolin and are not appropriate for stoma care as they prevent the skin barrier from sticking well to the skin.**
3. Rinse the area well and pat it dry with a soft towel

4. Any adhesive residue can be removed with an ostomy adhesive remover. Make sure that you cleanse the area well with warm water afterwards as the remover may interfere with the adherence of your new pouching system.
5. Shave around the stoma if needed. Use a light dusting of ostomy powder as a dry lubricant for shaving. Make sure that you shave away from the stoma to avoid cutting the stoma accidentally. Gently brush the skin clean. **Do not use a commercial hair removal product.**
6. Check the condition of your skin to make sure that it is not sore. Skin irritation can be caused by:
 - Traumatic removal of the pouching system
 - Incorrect pouching system fit from weight gain or loss
 - Leakage of stool onto skin
 - Sensitivity to products
 - Trapped moisture
7. If your skin is clear, wipe skin protector around your stoma and keep it dry. This step is optional. **If you have skin sensitivities or if your skin is red and sore, do not use skin protector.**
8. If your skin is red and sore, dust it lightly with ostomy powder. Brush away excess powder. Do not use creams or lotions because these products can interfere with the adherence of the new pouching system. **If you have persistent skin irritation, contact the NSWOC nurse.**
9. For the first 6 to 8 weeks, measure your stoma size each time you change the pouching system. Use a stoma measuring guide provided in the product packaging. After the stoma stops shrinking, measure it when your weight changes. Weight gain or loss changes the size of the stoma.
10. Cut a hole in the centre of the skin barrier. The hole should be 3mm (1/8th inch) bigger than your stoma. It should also be the same shape as your stoma. **Note:** Some skin barriers are available with pre-cut stomal openings. The center opening has already been created for your stoma size. They are simpler to use since there is no cutting involved. These skin barriers are not appropriate while your stoma is still swollen or if you have an oval or irregular-shaped stoma.

11. Remove the paper backing from the skin barrier.
12. If you choose to use ostomy paste, put a thin layer around the hole for the stoma.
13. If your skin has wrinkles, pull your skin slightly to make sure it is smooth before applying the skin barrier. Place the skin barrier against your abdomen, fitting it around your stoma. Gently press it into your skin.
14. Smooth the skin barrier with your fingertips to make sure it sticks to your skin.
15. Attach the pouch to the skin barrier. Give the pouch a firm tug to make sure it is secure.
16. Place your hand over the pouching system for 2 or 3 minutes. The warmth from your hand will help the skin barrier to stick.
17. Fasten the clamp on the pouch, or roll the outlet.
18. Secure your ostomy belt if you use one.

You may want to try a pouching system that is different from the one introduced to you by the NSWOC nurse. Ask the NSWOC nurse to show you samples or you can contact the manufacturers directly (see **Helpful resources**). They all have websites and toll-free numbers you can access for free samples. You are encouraged to try different brands and types of pouching systems. You can also attend open houses sponsored by various pharmacies to talk directly with ostomy sales representatives.

Note the features you prefer about pouching systems:

What about food?

As you recover from your operation, you will progress from clear fluids to a regular diet. There is very little or no change in your diet when you have a colostomy. **You do not need to be on a special diet** unless you have another medical condition or as advised by your doctor. Canada's Food Guide (see **Helpful resources**) recommends that you include a variety of healthy foods each day. The food guide includes the following recommendations:

- Have plenty of vegetables and fruits
- Eat protein-rich foods
- Choose whole grain foods
- Make water your drink of choice

Maintain a healthy weight. Eat 3 or more meals per day to help your colostomy function regularly. Drink fluids with meals.

You can enjoy alcoholic beverages in moderation. Keep in mind that drinking alcohol may not be advisable when taking prescribed medications. Seek the advice of your pharmacist.

Some foods can change the colour of the stool. Beets can turn the stool red. Broccoli, asparagus and spinach can darken the stool.

Eat in a pleasant, relaxed atmosphere.

The following guidelines can assist you in controlling some conditions commonly experienced by people with a colostomy:

Excessive gas

You may have more gas for the first few weeks after surgery. Slowly progress your diet. Tolerance to foods usually improves over time. Test new foods by adding them one at a time. Keep a record of foods that bother you. Try the problem food again at a later time.

If you have problems with excess gas, limit your intake of gas-forming foods:

- Dried peas and beans, lentils
- Melons
- Eggs
- Beer and carbonated drinks
- Onions, garlic and related vegetables
- Strong cheeses such as Roquefort
- Vegetables in the cabbage family such as broccoli, Brussels sprouts, cabbage, or cauliflower

Eat small amounts of well-cooked food. Do not eat large amounts of fresh fruits and vegetables, or foods that bothered you in the past.

At first, try only a small amount of the foods that produce gas. Increase portions one at a time.

Skipping meals may increase gas. Eating at regular intervals promotes regular bowel habits.

Chew food slowly and completely. Chew with your mouth closed and do not talk while eating. Chewing gum or using a drinking straw can increase gas.

Odour

While your pouching system is odour-proof, you may choose to limit your intake of foods that promote odour. Enjoy them at times when you are less likely to be concerned about odour.

Foods that cause odour are:

- Fish
- Eggs
- Onions, garlic
- Dried peas, beans, lentils
- Asparagus
- Fried foods
- Vegetables in the cabbage family, including turnip, cauliflower, Brussels sprouts, broccoli, cabbage
- Alcohol
- Strong cheeses
- Some spices

Foods that may be included in your diet to control odour are:

- Parsley
- Yogurt
- Buttermilk
- Cranberry juice

Diarrhea

It is normal for stool consistency to vary during the day depending on what you eat. When you have diarrhea, your stool is liquid or watery and you have more frequent bowel movements. You will need to empty your pouch more often. This section explains the effects of foods and changes that can be considered to prevent diarrhea.

Limit foods that may promote loose stools:

- Spicy foods
- Alcohol
- Coffee
- Salads and leafy vegetable
- Raw fruits and vegetables
- Natural laxatives such as caffeine, prunes, figs, licorice, and chocolate
- High fat foods
- Nuts and seeds
- High sugar foods and drinks

Try drinking fluids half an hour after meals, rather than with meals. Eating larger meals during the day and a smaller meal in the evening helps to reduce the stool volume at night.

Try foods that tend to thicken your stool:

- Applesauce (unsweetened)
- Bananas
- Cheese
- Soda crackers
- Toasted white bread
- Rice
- Marshmallows
- Creamy peanut butter
- Pretzels
- Bagels
- Pasta
- Tapioca
- Buttermilk
- Yogurt
- Boiled milk
- Porridges: oatmeal, oat bran, cream of wheat, cream of rice
- Potatoes

Constipation

You can still become constipated with a colostomy especially with regular use of some pain medications. To prevent constipation, eat foods that are high in fibre:

- Fresh fruits and vegetables
- Whole grain breads and cereals
- Foods with bran, nuts or seeds

Drink at least 6 to 8 glasses (1.5 to 2.0L) of fluids per day.

If you find you are passing hard stools, you may choose to eat stewed prunes or drink a glass of prune juice daily.

If the constipation persists, your doctor may recommend taking a stool softener or laxative. Contact your doctor for guidance.

What to wear?

Choose comfortable clothes to wear home from the hospital. Your pouching system is not visible when worn beneath your clothes and when emptied frequently.

You can wear the pouch over or under your underwear. Wearing pantyhose or a light girdle over your pouching system will not harm the stoma. Wearing snug underclothing over your stoma will help muffle the noise when passing gas and may help you feel more secure.

You can wear a belt or clothing with a waistband. Avoid a tight belt directly over the stoma. Make sure you can pass your fingers easily under your belt or waistband to make sure it is not too tight.

Pants that are worn tightly over the stoma won't allow the stool to enter the pouch and could cause harm to the stoma. Remember the stoma has no nerve endings so you can't feel it and won't know if you are injuring it.

If your colostomy is located above your beltline, you may wear an undershirt or camisole to keep the pouch secure and smooth. Wear loose outer clothing such as vests, sweaters, scarves and jackets help to conceal your pouching system.

When choosing swimwear, select dark colored suit with pattern to disguise your pouching system.

- For women, choose a suit with a small ruffle, skirt or panel.
- For men, choose boxer trunks with a high cut waist or longer leg.

If you are concerned that your pouching system is noticeable or that your pouch may shift when your suit is wet, wear a support garment or bike shorts under your suit. Customized swimwear with inside pockets is also available for additional pouch support (see [Helpful resources](#)).

How to bathe and shower

You can bathe or shower without taking off your pouching system. Dry both sides of the pouch to avoid skin irritation from trapped moisture.

Once your incision has healed completely, you can bathe or shower when your pouching system is off. This is the best way to clean your stoma and skin.

- Water does not harm or get into your stoma.
- Direct the forceful water stream away from the stoma.
- Do not use bath oils or soaps that leave an oily film on your skin. These can prevent your pouching system from sticking properly to your skin.
- After your bath or shower, dry well and place the new pouching system over your stoma.

What about work?

You need time to recover and learn to care for your colostomy after your operation. It can take several weeks to regain your former strength. As your recovery progresses, you will have many questions about your work. After a time of healing (approximately 4-6 weeks), you can slowly return to work.

Depending on the kind of work, you may need to make a few changes. Jobs that demand a lot of physical activity may have to be limited. You may find a gradual schedule will work best for you. Your surgeon and NSWOC nurse help you with this decision.

Keep extra supplies at your workplace in case you need them.

What about travel?

You can travel anywhere you like. You may start to drive when you are no longer taking opioid pain medications that affect your driving. Of course, you need to feel comfortable with your ileostomy care before you go to far-away places.

Here are a few tips to make your travels a success:

- Carry clean supplies wherever you go. You can carry supplies in a small cosmetic purse, waist pouch or briefcase.
- Be sure to take extra supplies. Take at least twice the supplies you normally use.
- When flying, bring supplies in your carry-on baggage so that you always have them with you in case your luggage is lost.
- If you are planning air travel, purchase pre-cut skin barriers so you will not need to carry scissors in your carry-on bag. Air pressure in airplanes will not affect your pouch.
- Keep your supplies out of the sun. Do not keep them in a hot place such as the trunk or glove compartment of a car.
- Wear your seatbelt when travelling in a car. Make sure that the belt goes across your lower abdomen. A soft pad can be placed under the seatbelt to protect the stoma.
- Drink bottled water if there is any doubt as to the safety of the water.
- Ask the NSWOC nurse or contact the Ostomy Canada Society about available resources in the area you plan to visit (see **Helpful resources**).

What about physical activity?

The day of your surgery, you may be visited by the physiotherapist who will begin working with you on your physical activity. They will work with you to help you return to your pre-operative mobility. Typically, the goal will be to walk to the bathroom and in the hallway with little to no assistance in order to be ready to go home. If your home is equipped with stairs, the physiotherapist will also ensure you are able to safely walk up and down the stairs before being discharged.

How to get out of bed after surgery

- Step one: bend both knees with feet on the bed, reach both arms towards the bedrail and in the direction of exit.
- Step two: push with your feet and roll onto your side.
- Step three: push yourself up using both arms on the bed. This reduces strain on the abdominal muscles, and protects the incision (reduces hernia risk).

It is normal to tire easily after your operation. Your body is using a great deal of energy to heal itself. Listen to your body and rest when you feel tired. Activity also helps with the healing. Don't be afraid to be active when you feel up to it. You can resume most normal activities once your pain is manageable, including sexual intercourse. Slowly increase your activity to your usual level. Follow a regular exercise program with activities like walking and bicycling. Exercise is good for you. Be as active as you like.

Temporary physical limitations

During the first 4-6 weeks after surgery while your body is still recovering, practice the following limitations:

- **Do not lift more than 10 to 15 pounds** (one laundry bin or 2 small bags of groceries) for the first 4-6 weeks after surgery.
- Do not do abdominal exercises, high intensity aerobic activities or weight training for 4-6 weeks after surgery.
- Avoid playing golf or any sport where there are twisting movements during this time.
- Avoid any sport where there is body contact during this time.

Ask your friends and family to help you with getting meals ready, grocery shopping, house cleaning and laundry.

Returning/starting a new sport

When your return to a sport, or try a new sport, remember the following:

- Ask your doctor about any limitations.
- Choose sports that are relaxing, comfortable and fun.
- Take it easy at first until you build up your strength. If you want to work out in a gym, avoid abdominal exercises for a while. Only introduce these gently, in gradual amounts after your surgeon says it is ok.
- Wear an ostomy belt, stoma shield, panty girdle or close-fitting briefs, if necessary.
- Talk to your NSWOC about products to meet your needs.

When returning to playing sports, protect your stoma from physical blows, from too tight clothing and from rigid equipment. Wear a stoma guard and if desired, an abdominal support as well. If you perspire a lot, wear an ostomy belt or apply extra tape around the edges of your pouching system to keep it secure.

Swimming and other water activities

Swimming, hot tubbing and saunas may be resumed once your incision heals completely. If you enjoy water sports, try these tips:

- Don't go swimming right after putting on a new pouching system.
- Make sure your pouch is empty or wear a stoma cap or mini-pouch instead of a standard sized pouch.
- Cover the edges of your pouching system with waterproof tape or seal.
- You may choose to change and towel off in a bathroom stall for more privacy.
- You may choose to put on a dry, oversized T-shirt as a cover-up while you change.
- You may choose to wear your bathing suit under your clothing to avoid having to change.
- Arrange to go swimming when your colostomy is less likely to be active (for example: during an empty stomach, not immediately after eating).
- A dry suit is easier to take off than a wet one.

Hernia prevention

The abdominal muscles need time to heal and become stronger after surgery. In addition to avoiding heavy lifting, intensive abdominal exercises should begin no sooner than 3 months after your surgery. Always wait until your surgeon says it is okay to resume abdominal exercises. Maintaining a good posture will help to develop your muscles. Your weight should be kept within the appropriate Body Mass Index (BMI) range for your height. Supporting your stoma and abdomen while coughing in the first few months after surgery will also assist in reducing the chance of developing a herniation. Gentle abdominal exercises you can try after surgery once your wound has completely healed include:

- **Pelvic tilting:** Lie on your back on a firm surface with knees bent and feet flat. Pull your tummy in, tilt your bottom upwards while pressing the middle of your back into the surface (floor/bed) and hold for 2 seconds, let go slowly. Repeat 10 times daily.
- **Knee rolling:** Lie on your back on a firm surface with knees bent and feet flat on the surface (floor/bed). Pull your tummy in and keeping your knees together, slowly roll them from side to side. Repeat 10 times daily.
- **Partial abdominal sit ups:** Lie on your back on a firm surface with knees bent and feet flat on the surface (floor/bed). Place your hands on the front of your thighs and pull your tummy in. Lift your head off the pillow. Do not roll your back up off of the surface. Hold for 3 seconds, then slowly return to starting position. Repeat 10 times daily.

What about intimacy?

Your colostomy does not need to limit your sexual activity. It takes time for both partners to understand and get used to each other's feelings. Open discussion with your spouse or partner helps in your adjustment to changes in your body. Your surgeon will talk to you about any sexual problems you might have when you resume sexual activity.

If the rectum has been removed, men sometimes have nerve damage in their lower abdomen that makes it difficult to have an erection. This damage may go away after a few months or it may last. If this is a problem for you, see your doctor. Treatments are available.

If the rectum had been removed, some women find the position of their vagina changes. This can make it difficult to have a climax. Try different positions to find what works best for you.

Pregnancy may be a concern when sexual activity is resumed. Consult your physician to discuss this. Many women with a colostomy have had successful pregnancies and vaginal deliveries. If you do become pregnant, consult your NSWOC nurse since your pouching system may require some adjustments as the abdomen increases in size.

Sometimes, the problem is emotional. It helps to make sure your pouching system is clean and secure. Stoma caps or mini-pouches can be used as these are much smaller than a standard pouch and are discreet. Also, try making your pouching system less noticeable by wearing customized underwear or lingerie (see [Helpful resources](#)).

Only through time and experience will you know how it will be for you. If you do have problems, talk with your surgeon or NSWOC nurse. They can help you find solutions or direct you to other services.

Medications

Most medications can be taken safely when you have a colostomy. Most medications are absorbed in the stomach and small bowel. Keep in mind that certain medications such as vitamins and antibiotics can affect the odour, colour and consistency of the stool.

Regular use of pain medications can result in severe constipation. To prevent constipation, refer to the [Constipation](#) section.

Discuss any concerns with your pharmacist or doctor.

Adjusting to your colostomy

Your feelings about your colostomy may change from day to day. Sometimes you may feel confused, angry, or sad. Other times you may feel happy that the operation is over. It is natural to feel this way after your body has had such major change. It helps to talk about your feelings.

As you get used to having a colostomy and feel sure about your care, you will find it easier. This change begins in the hospital and continues when you are at home. Each new experience you master helps you feel more comfortable and confident. This takes time, so be patient with yourself.

Not everyone needs to know that you have a colostomy. You can decide whom to talk to about your colostomy. It helps to talk to your family, close friends or others who have a colostomy. Discussion with a volunteer from a local chapter of the Ostomy Canada Society who has experienced similar surgery is helpful (see [Helpful resources](#)).

Your family and close friends also need time to understand and accept your colostomy. You can help them feel more relaxed about your colostomy if you feel at ease, and treat it as a part of you. In time, they will see that you are still the same person that they have always known. They will see that you can return to your previous family and social roles.

Before you go home

You will be required to demonstrate your independence with pouch care before going home.

A Home and Community Care Case manager (or Local Health Integration Network –LHIN-manager) calls you in the hospital to:

- Talk to you about your community care needs
- Arrange for a nurse to help you with care
- Arrange for other health care that you may need
- Determine if you qualify for supplies

The Ontario government pays for Home and Community Care. It is a service for Ontario residents who have a valid Ontario Health Insurance Plan (OHIP).

Post-surgical follow-up

Within one week after discharge, you will be encouraged to participate with your pouching system change. Your visiting nurse will support you and encourage you to become independent with pouching system changes.

You will receive an appointment for follow-up with your surgeon 4 to 6 weeks after your hospital discharge. The NSWOC from Hotel Dieu Hospital site (HDH) will be able to assist you with any ostomy or skin related difficulties that may arise after you have gone home.

The HDH site NSWOC nurse will also be available to see you in the clinic for reassessment of your colostomy at any point in your ostomy journey. Bring along your supplies for a pouching system change. Pre-booked visits are arranged between the NSWOC and the patient by calling the ostomy clinic (refer to page 37).



How to pay for supplies

If you are an Ontario resident and have a colostomy for longer than 6 months, you can apply for a grant from the **Assistive Devices Program (ADP)**. Your surgeon or family doctor must certify that you have a colostomy and sign the application form. Through this program, you are eligible for \$975.00 per year. If you receive social assistance benefits, you are eligible to receive \$1,300.00 per year. If you are a resident of a Long-Term Care facility, you are eligible for \$1,300.00 per year. You receive two cheques from ADP, in equal amounts, six months apart.

You can also set up for direct deposit into your bank account.

You must keep your receipts or copies of your receipts for two years. The Ministry of Health may ask you to produce these receipts to confirm your claim.

It is important to note that these benefits may not fully cover the annual costs of your colostomy.

For income tax purposes, you can claim receipts for costs that exceed the amount of your ADP grant. For more information, talk to your NSWOC nurse or contact ADP:

Ministry of Health and Long-Term Care
Assistive Devices Program
5700 Yonge Street, 7th Floor
Toronto, ON
M2M 4K5
Toll free 1-800-268-6021
TDD (for the hearing impaired) 1-800-387-5559
www.health.gov.on.ca

If you have a **private medical insurance** plan, contact your agent. Many private insurance plans will reimburse you for some of the costs of your ostomy supplies. If you have ADP funding, your private insurance may cover some of the costs that exceed the ADP funding.

If you have an ostomy for a continuous period for at least 12 months, you may be eligible for the **Disability Tax Credit**. This is a non-refundable tax credit used to reduce the amount of tax payable on your income. Your physician must certify the tax credit certificate and it must also be approved by Canada Revenue Agency (CRA). A T2201 Form is required and can be downloaded from the CRA website or picked up from your local Service Canada Centre office.

For more information, talk to your NSWOC nurse or contact CRA: <http://www.cra-arc.gc.ca/>

What to do if...

Stoma

Stoma turns dark purple or black

Go to the emergency room/urgent care centre.

Stoma bulges out from your abdomen

Call your surgeon or go to the emergency room/urgent care centre.

Stoma opening becomes narrow and bowel movements are hard to pass

Call the NSWOC nurse.

Remember that it takes 6 to 8 weeks for your stoma to shrink to its final size.

Excessive bleeding

A small amount of bleeding is normal.

If a lot of blood comes from the stoma, call your surgeon or go to the Emergency Department.

Stoma becomes hidden in abdomen

Call the NSWOC nurse.

Skin

Bruising

Call the NSWOC nurse if the bruising does not go away after 4 weeks.

Remember that it may take longer if you take blood thinners.

Rash

Do not use skin protector.

Remove the pouching system and wash your skin in the shower or bath. Pat your skin dry. Apply a light dusting of ostomy powder. Place a new pouching system.

If the rash does not get better, call the NSWOC nurse.

Pouching system

- Leaks** Use soap without oil to wash the skin around your stoma before placing a new pouching system.
- Make sure that your skin is dry before placing a new pouching system.
- Use a hair dryer on a low setting to blow the skin dry.
- Spread a thin layer of ostomy paste around the opening on the back of the skin barrier to help create a good seal.
- Place the pouching system on a smooth skin surface. Pull gently on your skin to remove wrinkles and creases.
- Weight gain or loss changes the fit of your pouching system. Call the NSWOC nurse for re-fitting.
- Painful to remove* *Add adhesive removers to your routine.*
- Shave the hair from the skin around your stoma before placing a new pouching system.*

Call the NSWOC nurse if any problems with your stoma, skin or pouching system persist. Refer to Page 37 for contact numbers.

Note your stoma/skin/pouching system concerns

About your colostomy closure

Some colostomies can be closed (reversed). The bowels need time to heal before your temporary colostomy is closed. Your surgeon will tell you when the colostomy is ready to be closed. This usually takes place within 3-6 months.

You prepare the same way that you did for the colostomy surgery (see **Before your operation**).

The surgeon usually operates through the same incision or at the stoma site on your abdomen. After the operation, a dressing covers the former colostomy site.

After your operation, you may be allowed to eat and drink 2 hours after your operation.

About 2 to 3 days after your operation, you should be discharged from hospital.

Information to remember

Name of operation:

Date of operation:

Discharge date:

Surgeon:

telephone: _____ ext: _____

NSWOC nurse:

KGH site: _____

KGH site: _____

HDH site: _____

Ostomy clinic location:

Hotel Dieu Hospital (Jeanne Mance 5)
Telephone: 613-544-3400 ext. 3720

Type of colostomy:

- loop end other
-
- temporary permanent

Stoma size:

Stoma shape: _____

Date of last pouching system change:

Pouching system type:

- 1-piece system 2 piece system

Helpful resources

Canada's Food Guide

www.food-guide.canada.ca

Canada Revenue Agency

Disability tax Credit

www.cra-arc.gc.ca

Canadian Cancer Society

National Office

55 St. Clair Avenue West Suite 500

Toronto, ON M4V 2Y7

Toll Free: 1-888-939-3333

www.cancer.ca

Canadian MedicAlert® Foundation Inc.

Morneau Shepell Centre II

895 Don Mills Road, Suite 600

Toronto, ON M3L 1W3

Toll Free: 1-800-668-1507

www.medicalert.ca

Cancer Care Ontario

620 University Avenue, 15th floor

Toronto, ON M5G 2L7

Toll Free: 1-888-939-3333

www.cancercareontario.ca

Colorectal Cancer Association of Canada

4576 Yonge Street, Suite 608

North York, ON M2N 6N4

Toll Free: 1-877-50colon (26566)

www.colorectal-cancer.ca

Coloplast Canada Corporation

2401 Bristol Circle, Suite A205

Oakville, ON L6H 5S9

Toll Free: 1-866-293-6349

www.coloplast.ca

ConvaTec

Suite 250, 1425 Trans-Canada Hwy.

Dorval, QC H9P 2V3

Toll Free: 1-800-465-6302

www.convatec.ca/CA/en (English)

www.convatec.ca/CA/fr (French)

Crohn's and Colitis Canada

Suite 600, 60 St. Clair Avenue East

Toronto, ON M4T 1N5

Toll Free: 1-800-387-1479

www.crohnsandcolitis.ca

**Friends of Ostomates Worldwide
(F.O.W Canada)**

301 Benlock Road
Grafton, ON K0K 2G0
Phone: 647-951-3940
www.fowc.ca

Hollister Limited

95 Mary Street
Aurora, ON L4G 1G3
Toll Free: 1-800-263-7400
www.hollister.ca

**Nurses Specialized in Wound, Ostomy,
Continence Canada**

66 Leopolds Drive
Ottawa, ON K1V 7E3
Toll Free: 1-888-739-5072
Fax: 1-514-739-3035
www.nswoc.ca

Ostomy Canada Society

Suite 210, 5800 Ambler Drive
Mississauga, ON L4W 4J4
Phone: 905-212-7111
Toll Free: 1-888-969-9698
www.ostomycanada.ca

Stealth Belt

210 West Oakland Avenue Suite B
Johnson City, TN 37604
Toll Free: 1-800-237-4491
www.stealthbelt.com

Weir Comfees

P.O Box 255 Alliston, ON L9R 1V5
Toll Free 1-866-856-5088
www.weirdcomfees.com

Additional resources

Glossary

Anus	opening at the lower end of the rectum through which stool is passed during a bowel movement
Bowel	the part of the digestive system between the stomach and the anus where food is broken down and stool is stored. This is also known as the intestines.
Bowel Movement	when stool is eliminated from the colostomy or rectum
Bowel Obstruction	when the bowel is blocked, either partially or completely, and the stool cannot pass through
Cancer	a disease involving abnormal cell growth with the potential to invade or spread to other parts of the body and to destroy normal tissue
Catheter	a thin, soft tube used to drain fluids such as urine
Colon	the last 5 feet of the digestive system, also called the large bowel
Colostomy	a surgical opening through the abdomen into the colon
Constipation	stool becomes hard; bowel movements are infrequent and difficult to pass
Diarrhea	frequent, watery bowel movements
Diverticulitis	inflammation of a small sac in the lining of the large bowel
Drain	a thin, flexible tube used to drain fluids out of the body
Enema	fluid to put into the rectum to clean out stool
Esophagus	a muscular tube that carries food or fluids from your throat to the stomach
Incision	the cut, or wound, made by the doctor during the operation

Inflammatory Bowel Disease	term for diseases that result in prolonged inflammation of the bowel such as ulcerative colitis and Crohn's
Large Bowel	the last 5 feet of the digestive system, also called the colon
Laxative	medication that cause bowel movements
Mucus	a clear jelly-like substance made by the lining of the digestive tract to keep it moist
Nurse Specialized in Wound, Ostomy, Contenance	a nurse with special training in caring for persons with ostomies; also known as an NSWOC nurse
Ostomy	a surgical opening into the bowel, a short form for colostomy
Pouching System	the pouch worn over your stoma to collect stool
Rectum	the end part of the large bowel that stores stool
Skin barrier	the part of the pouching system that adheres to the abdomen and protect the skin around the stoma
Small bowel	the part of the digestive system between the stomach and large bowel, that breaks down the food and absorbs nutrients (also called the small intestine)
Stoma	an opening through the abdomen to allow the bowel pass stool, also the Greek word for "mouth"
Stool	the waste product of digestion, also called a bowel movement

Transforming care, together™

Kingston Health
Sciences Centre

Centre des sciences de
la santé de Kingston



Hôpital
Hotel Dieu
Hospital



Hôpital Général de
Kingston General
Hospital

76 Stuart St. Kingston, ON, Canada K7L 2V7

Tel: 613-549-6666 www.KingstonHSC.ca

Aussi disponible en français

KHSC Stores 359

December 2020